

Codex[®]

BVMO Panel

Screening Protocol

Document # PRO-013-04

Reactions of Interest

Bayer-Villiger Oxidation



Sulfoxidation



Codex[®] BVMO Screening Kit General Information

- The Codex[®] BVMO Panel is a collection of Baeyer-Villiger Monooxygenases (BVMOs) in 96-well plates containing 100 µL BVMO lysate in each well. BVMOs have been developed for the asymmetric oxidation of sulfide to sulfoxide. The enzymes included in this panel were developed using Codexis' CodeEvolver[®] protein engineering technology and have been engineered for improved activity and stability.
- Recommended storage temperature for the plates is -20 °C.

Screening Reagents Required

- Codex[®] BVMO Panel. Two 96-well plates, provided.
- 200 mg recycling KRED, provided.
- 100 mg NADP⁺, provided.
- 200 mg GDH-105, provided.
- 50 or 100 mM potassium phosphate buffer, pH 9.0, not provided.
- Isopropanol or glucose, not provided.

Note: If the substrate contains a ketone or aldehyde function that reacts with the KRED, then consider replacing the KRED recycling system with GDH/glucose recycling system.

Codex® BVMO Panel

Codex® BVMO Screening Procedure

1. Defrost plates, which contain 100 μ L lysate per well, at 4 °C. Centrifuge (4000 rpm, 2 min, 4 °C).
2. Setup the assay using either the KRED/IPA or the GDH/glucose cofactor recycling system per the table below for a final reaction volume of 300 μ L per well.

Reagent concentration using KRED/IPA for cofactor recycling	Volume per well	Volume per plate	Final concentration
Lysate	100 μ L	9.60 mL	33% v/v
Solution A 1.8 g/L KRED-P1-A12 0.9 g/L NADP+ 50 mM buffer	170 μ L	16.32 mL	57% v/v 1 g/L KRED-P1-A12 0.5 g/L NADP+
Solution B 10–50 mg/mL substrate in Isopropanol	30 μ L	2.88 mL	10% v/v 1–5 mg/mL substrate 10% isopropanol
Total Volume	300 μ L	28.8 mL	

Reagent concentration using GDH/glucose for cofactor recycling	Volume per well	Volume per plate	Final concentration
Lysate	100 μ L	9.60 mL	33% v/v
Solution A 1.8 g/L GDH-105 0.9 g/L NADP+ 100 mM buffer	100 μ L	9.60 mL	33% v/v
Solution B 10–50 mg/mL substrate in buffer or co-solvent 15–75 mg/mL glucose	30 μ L	2.88 mL	10% v/v 1–5 mg/mL substrate 1.5–7.5 mg/mL glucose
Total Volume	300 μ L	28.8 mL	

3. Assay setup:
 - a. Unseal the plates.
 - b. Add 170 μ L of Solution A to each well.
 - c. Add 30 μ L of Solution B to each well.
 - d. Seal the assay plate using a plate sealer (180 °C, 3 sec).
4. Shake the plates on titer plate shaker at low speed and at room temperature for 24 hours.

Codex® BVMO Panel

Codex® BVMO Panel Work Up and Analysis

1. Reaction quenching:
 - a. Centrifuge the assay plates (4000 rpm, 1 min, 4 °C).
 - b. Unseal the assay plates.
 - c. Add 1000 µL of quenching solvent (e.g. acetonitrile, methanol, or MTBE) to each well.
 - d. Seal the quenched plates (180 °C, 3 sec).
2. Shake the plates on titer plate shaker (medium-high speed) at room temperature for 20 minutes.
3. HPLC sample preparation:
 - a. Centrifuge the assay plates (4000 rpm, 20 min, 4 °C).
 - b. Unseal the quenched plates.
 - c. Transfer 200 µL of samples from quenched plates to round-bottom shallow well plates (Corning 3365).
 - d. Seal the shallow well plates (180°C, 2 sec).
4. Analyze using HPLC method of choice.

For further information,
please contact us at sales@codexis.com